What is it?

Bell's Palsy is a condition where there is involvement of the facial nerves, causing a droop on one side of the face. A picture of the facial nerve is on the front of this pamphlet for reference.


This can present similarly to a stroke, but is not caused by a stroke. Full recovery can take an upwards of 6 months.



Stroke Prevention Clinic

Saskatoon:

P: 306-655-6802 | F: 306-655-6803 Regina: P:306-766-3959 | F: 306-766-3968



For More Information

www.heartandstroke.ca

Mayo Clinic. (2022, May 4). *Bell's Palsy*. https:// www.mayoclinic.org/diseases-conditions/bells-palsy/ symptoms-causes/syc-20370028


Cleveland Clinic. (2023, August 14). Bell's Palsy. https:// my.clevelandclinic.org/health/diseases/5457-bells-palsy





Bell's Palsy

Stroke Prevention Clinic



Used with permission of Mayo Foundation for Medical Education and Research, all rights reserved.



saskhealthauthority.ca

Healthy People, Healthy Saskatchewan

Symptoms

Symptoms of Bell's Palsy include:

- facial weakness on one side of the face
- facial droop
- drooling
- loss of sense of taste
- jaw or ear pain
- nnability to close eyelid on affected side

Causes

Bell's Palsy most often has an unknown cause. It is believed to be due to the inflammation and swelling of the nerve that controls that side of the face. It can also occur after a viral infection, such as a cold sore

Diagnosis

Based on your symptoms and neurological exam, your neurologist can determine whether you have Bell's palsy. Sometimes, imaging may be ordered to rule out other causes that can mimic these symptoms. Bloodwork may also be done to determine if you have any current viral infections.




Image provided from SHA Creative Services. Facial nerve paralysis, Bell's palsy, illustration showing female with one sided facial nerve paralysis. AdobeStock; 2021 [cited 2024 APR 04]. Available from: <u>https://stock.adobe.com/ca/images/facial-nerve-paralysis-bell-s-palsyillustration-showing-female-with-one-sided-facial-nerveparalysis/424121568</u>

Treatment

Treatment for Bell's palsy includes:

Medications:

- Corticosteroids an antiinflammatory medication like prednisone can help reduce swelling
- Antiviral drugs if the cause is viral (cold sores)
- Eye drops and eye ointment will also be used due to the inability to keep the eyelid closed. In some cases, an eye patch may be used to protect the affected eye.

A physiotherapist can show you how to massage and exercise the facial muscles in severe cases.

If at any time you are concerned about your symptoms, or they are unusual for you, please seek medical attention.

The Saskatchewan Health Authority works in the spirit of truth and reconciliation, acknowledging Saskatchewan as the traditional territory of First Nations and Métis People.

CS-PIER-0105

April 2024